

Lanhuong Thi Pham, D.P.M.

Joel Robinson, D.P.M.

119 Medical Circle Sulphur Springs, Texas 75482 including Satellite offices

P: (903) 885-2754 **F:** (903) 347-1207

Email: phamfootdoc@gmail.com

NAIL MATRIXECTOMY - POST PROCEDURE CARE

SUPPLIES

- 1. Hydrogen Peroxide
- 2. 1-inch Band Aids
- 3. Q-Tips
- 4. Antibiotic Ointment* (Neosporin or Triple Antibiotic)
- 5. Pain Management OTC Medication
 - a. Advil (Ibuprofen)
 - b. Aleeve (Naproxen)
 - c. Tylenol (Acetaminophen)
- 6. Epsom Salt

INSTRUCTION

- 1. Keep the procedure site and the bandage dry, stable and intact for 24 Hours a day. To remove the dressing, soak it in Salt water to make it easier to release from the toe.
- 2. Soak feet on the SECOND day by adding 2 table spoons Epsom Salt to 1 quart of warm water. Soak it for 10-20 minutes.
- 3. Remove the bandage and clean the sides of the nail border with Hydrogen Peroxide using the Q-Tips.
- 4. Dry the toe. Apply Antibiotic ointment* & then apply the band aid lightly over the area.
- 5. Repeat steps #2, #3 & #4 daily for TWO weeks.
- 6. It is normal for the wound to drain a CLEAR liquid for a few days. DO NOT worry if you see a little drainage from the wound.
- 7. For pain management, please take over the counter Advil, Aleeve, Tylenol or medication prescribed by the doctor as directed.
- 8. If there is **REDNESS** coming from the toe towards the foot or **RED STREAKS** past the knuckle area of the toe, please contact the doctor's office

* Patients with allergies to Neosporin may use Bacitracin or Polysporin instead